

University of Groningen

Element specific magnetic moments of gadolinium iron garnet probed by soft-x-ray magnetic circular dichroism

Rudolf, P.; Sette, F.; Tjeng, L.H.; Meigs, G.; Chen, C.T.

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1991

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rudolf, P., Sette, F., Tjeng, L. H., Meigs, G., & Chen, C. T. (1991). *Element specific magnetic moments of gadolinium iron garnet probed by soft-x-ray magnetic circular dichroism*.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Element specific magnetic moments of gadolinium iron garnet probed by soft-x-ray magnetic circular dichroism (abstract)

P. Rudolf, F. Sette, L. H. Tjeng, G. Meigs, and C. T. Chen
AT&T Bell Laboratories, Murray Hill, New Jersey 07974

The element specific magnetic moments of Gd and Fe in a gadolinium iron garnet ($\text{Gd}_3\text{Fe}_5\text{O}_{12}$) were probed at various temperatures by measuring the soft-x-ray magnetic circular dichroism (SXMCD) at the Gd $M_{4,5}$ and the Fe $L_{2,3}$ adsorption edges in an external field of 5 kG. The soft-x-ray absorption spectra were recorded with the photocurrent method using highly circularly polarized soft x rays recently obtained at the AT&T Bell Laboratories Dragon synchrotron radiation beamline.¹ The temperature dependence of the SXMCD signal at each edge allows one to independently determine the magnetic ordering of each specific ion. Our individually probed magnetic moment of the Gd^{3+} ion and of the two Fe^{3+} ions in different sites confirm the earlier calculations which predict a reversal of the local magnetic orientation at ~ 280 K. This work demonstrates that SXMCD is a valuable tool for probing the element and site specific magnetic properties of multicomponent systems. The applications of this novel technique to the investigation of transition metals and rare earths in ferromagnetic alloys, compounds, disordered, and impurity systems will be addressed.

¹C. T. Chen, F. Sette, Y. Ma, and S. Modesti, *Phys. Rev. B* **42**, 7262 (1990).

²R. Pauthenet, *J. Appl. Phys.* **29**, 253 (1958).